

Final state strong interaction constraints on weak $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ decay amplitudes

B. Loiseau · J.-P. Dedonder · A. Furman ·
R. Kamiński · L. Leśniak

Received: date / Accepted: date

Abstract Weak decay tree and annihilation - t-channel W^- exchange amplitudes for the $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ process are calculated using quasi two-body QCD factorization approach and unitarity constraints. Final state strong $K\pi$ and $\pi\pi$ interactions in S , P and D waves are described through corresponding form factors including many resonances. Preliminary results compare well with the effective mass distributions of the Belle and BABAR Collaboration analyses.

Keywords QCD factorization · Final state interactions · Unitary form factors

PACS 13.25.Hw · 13.75.Lb

1 Introduction

Why should one study the weak decays $D^0 \rightarrow K_S^0 \pi^+ \pi^-$? First, the recent measurements of the D^0 - $\bar{D}^0$ mixing parameters for this self-conjugate reaction by the BABAR [1] and Belle [2] Collaborations could show the presence of new physics contribution beyond the standard model. Second, the Cabibbo-Kobayashi-Maskawa, CKM, angle γ can be evaluated from the analyses of the $B^\pm \rightarrow D^0 K^\pm$, $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Third, one can learn about the final state meson-meson interactions, the meson resonances decaying into different meson-meson pairs and their interferences in the Dalitz plot. One can also perform a partial wave analysis of decay amplitudes. In

Presented at the 20th International IUPAP Conference on Few-Body Problems in Physics, 20 - 25 August, 2012, Fukuoka, Japan

B. Loiseau, J.-P. Dedonder
Laboratoire de Physique Nucléaire et de Hautes Énergies, Groupe Théorie, Université Pierre et Marie Curie et Université Paris-Diderot, IN2P3 et CNRS, 4 place Jussieu, 75252 Paris, France
E-mail: loiseau@lpnhe.in2p3.fr

A. Furman
ul. Bronowicka 85/26, 30-091 Kraków, Poland

R. Kamiński, L. Leśniak
Division of Theoretical Physics, The Henryk Niewodniczański Institute of Nuclear Physics,
Polish Academy of Sciences, 31-342 Kraków, Poland

addition, constraints from quasi two-body QCD factorization [3], QCDF, approach will allow to test theoretical models of form factors entering in the decay amplitudes.

Quasi two-body factorization Following a program devoted to the understanding of the rare three-body B decays (see e.g. Ref. [4]) the presently available $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ data are analyzed in the framework of QCDF. Neglecting the small CP violation in K^0 decays, it will be assumed that $|K_S^0\rangle = (|K^0\rangle + |\bar{K}^0\rangle)/\sqrt{2}$. The three-meson final states $\bar{K}^0 \pi^+ \pi^-$ are approximated (quasi two-body approximation) as being formed by a meson-meson state, $[\bar{K}^0 \pi^+]_{S,P,D}$ or $[\bar{K}^0 \pi^-]_{S,P,D}$ or $[\pi^+ \pi^-]_{S,P,D}$ in a S, P or D wave created by a $q\bar{q}$ pair and the remaining meson, π^- , π^+ or $\bar{K}^0$, respectively. Amplitudes are derived from the weak effective Hamiltonian, H_{eff} which is a superposition of left-handed quark current-current operators. For instance one has the operator $O_1 = j_1 \otimes j_2$ where $j_1 = \bar{s}_\alpha \gamma^\nu (1 - \gamma^5) c_\alpha \equiv (\bar{s}c)_{V-A}$ and $j_2 = \bar{u}_\beta \gamma_\nu (1 - \gamma^5) d_\beta \equiv (\bar{u}d)_{V-A}$, α and β being color indices. Applying QCDF to the part of the amplitude proportional to the product of the CKM quark mixing matrix elements $V_{cs}^* V_{ud} \equiv A_1$ one has,

$$\begin{aligned} \langle \bar{K}^0 \pi^- \pi^+ | H_{eff} | D^0 \rangle &\simeq \frac{G_F}{\sqrt{2}} A_1 \left\{ a_1 \langle \pi^+ | j_2 | 0 \rangle \langle [\bar{K}^0 \pi^-]_{S,P,D} | j_1 | D^0 \rangle \right. \\ &+ a_2 \langle \bar{K}^0 | j'_2 | 0 \rangle \langle [\pi^+ \pi^-]_{S,P,D} | j'_1 | D^0 \rangle + a_2 \langle 0 | j'_1 | D^0 \rangle \langle [\bar{K}^0 \pi^-]_{S,P,D} \pi^+ | j'_2 | 0 \rangle \\ &\left. + \langle \bar{K}^0 | \pi^+ \pi^- \rangle_{S,P,D} | j'_2 | 0 \rangle \right\} \end{aligned} \quad (1)$$

where $G_F = 1.166 \times 10^{-5} \text{ GeV}^{-2}$ is the Fermi coupling constant, $a_{1,2}$ are effective QCD Wilson coefficients and $j'_1 = (\bar{u}c)_{V-A}$, $j'_2 = (\bar{s}d)_{V-A}$ derive from $j_{1,2}$ via a Fierz transformation. In Eq. (1) it appears that the cross product of bilinear quark currents in O_1 factorizes, after the introduction of the vacuum state, into the product of two matrix elements. The first ones are proportional to the π^+ , $\bar{K}^0$, D^0 decay constants f_π, f_K, f_{D^0} since $\langle \pi^+ | j_2 | 0 \rangle = if_\pi p_{\pi^+}$, $\langle \bar{K}^0 | j'_2 | 0 \rangle = if_K p_{K^0}$, $\langle 0 | j'_1 | D^0 \rangle = -if_{D^0} p_{D^0}$, p_{π^+}, p_{K^0} and p_{D^0} being the π^+ , $\bar{K}^0$ and D^0 four momenta, respectively. The second ones are transition matrix elements or form factors as with $M_i = \bar{K}^0, \pi^\pm$, $j = j_{1,2}$ or $j'_{1,2}$ one has, $\langle [M_1 M_2]_{S,P,D} | j | D^0 \rangle = \langle \bar{D}^0 [M_1 M_2]_{S,P,D} | j | 0 \rangle$ and $\langle [M_1 M_2]_{S,P,D} M_3 | j | 0 \rangle = \langle [M_1 M_2]_{S,P,D} | j | \bar{M}_3 \rangle$. It can be shown from field theory and using dispersion relations [5] that these form factors can be calculated exactly if one knows the D^0 - $[M_1 M_2]_{S,P,D}$ or M_3 - $[M_1 M_2]_{S,P,D}$ strong interactions at all energies.

2 Decay amplitudes

Different type of amplitudes Amplitudes with $c \rightarrow s\bar{u}d$ transition are proportional to $V_{cs}^* V_{ud}$ where $V_{cs} \approx V_{ud} \approx \cos \theta_C \approx 0.975$, θ_C being the Cabibbo angle. There are 7 such allowed tree amplitudes: 3 for the $\pi^+ [\bar{K}^0 \pi^-]_{S,P,D}$, 3 for the $\bar{K}^0 [\pi^+ \pi^-]_{S,P,D}$ and 1 for the $\bar{K}^0 \omega (\omega \rightarrow [\pi^+ \pi^-]_P$ by G -parity violation) final states. Amplitudes with $c \rightarrow d\bar{u}s$ transition, proportional to $\sin^2 \theta_C \approx (0.225)^2$, are doubly Cabibbo suppressed. There are 6 of them as the W meson cannot couple to the $[K^0 \pi^+]_D$ final state. There are also 7 allowed tree annihilation or t -channel W -exchange amplitudes corresponding to the $c\bar{u}$ annihilation into $s\bar{d}$ and 7 doubly Cabibbo suppressed annihilation amplitudes from the $c\bar{u}$ annihilation into $d\bar{s}$. Altogether, the quasi two-body QCDF approach leads to 27 non-zero - 13 tree and 14 annihilation - amplitudes,

Transition matrix elements Several meson resonances can decay into the two meson final states in S, P or D wave. For the kaon-pion subsystems, the scalar resonances, $K_0^*(800)^\pm$ or $\kappa^\pm$ and $K_0^*(1430)^\pm$ decay into $[\bar{K}^0\pi^\pm]_S$, the vector resonances, $K^*(892)^\pm$, $K_1(1410)^\pm$, $K^*(1680)^\pm$ into $[\bar{K}^0\pi^\pm]_P$ and the tensor resonances $K_2^*(1430)^\pm$ into $[\bar{K}^0\pi^\pm]_D$. For the pion-pion subsystems, the scalar $f_0(600)$ or σ , $f_0(980)$, $f_0(1400)$ decay into $[\pi^+\pi^-]_S$, the vector, $\rho(770)^0$, $\omega(782)$, $\rho(1450)^0$, $\rho(1700)^0$ into $[\pi^+\pi^-]_P$ and the tensor $f_2(1270)$ into $[\pi^+\pi^-]_D$. This leads to a rich interference pattern in the Dalitz plot. These resonances are used to write the three-meson transition to the vacuum as,

$$\langle M_1(p_1)[M_2(p_2)M_3(p_3)]_{S,P,D} | j'|0 \rangle \simeq G_{R_{S,P,D}^{M_2M_3}}(s_{23}) \langle M_1(p_1)R_{S,P,D}^{M_2M_3} | j'|0 \rangle, \quad (2)$$

where $s_{23} = (p_2 + p_3)^2$. The vertex functions $G_{R_{S,P,D}^{M_2M_3}}(s_{23})$ describe the $R_{S,P,D}^{M_2M_3}$ resonance decays into the states $[M_2M_3]_{S,P,D}$. A similar equation holds [6] for the D^0 transitions [see Eq. (1)] to two-meson states. As an example of application of Eq. (2) one can choose $M_1 \equiv \bar{K}^0$, $[M_2M_3]_S \equiv [\pi^+\pi^-]_S$ and $R_S^{\pi^+\pi^-} \equiv f_0(980)$, so that,

$$\langle \bar{K}^0(p_1)f_0(p_2+p_3)|(\bar{s}d)_{V-A}|0 \rangle = -i \frac{m_{K^0}^2 - s_{23}}{p_{D^0}^2} p_{D^0} F_0^{\bar{K}^0 f_0}(m_{D^0}^2) (m_{D^0}^2) + 1 \text{ term}, \quad (3)$$

where $p_{D^0} = p_1 + p_2 + p_3$. The $\bar{K}^0$ to f_0 scalar transition form factor, $F_0^{\bar{K}^0 f_0}(m_{D^0}^2)$, related to the $\bar{K}^0 f_0$ interaction at $(p_K + p_{f_0})^2 = m_{D^0}^2$, is a complex number to be fitted. The extra “1 term” gives a null contribution when multiplied, in Eq. 3, by $\langle 0 | (\bar{u}c)_{V-A} | D^0 \rangle$ [see Eq. (1)]. The vertex function $G_{f_0(980)}(s_{23}) \simeq \chi_2 F_0^{\pi^+\pi^-}(s_{23})$ where χ_2 is a constant fitted to data and $F_0^{\pi^+\pi^-}(s_{23})$ is the complex pion scalar form factor [4] related to $\langle [\pi^+\pi^-]_S | (\bar{u}u)_{V-A} | 0 \rangle$. The other vertex functions for the kaon-pion [7] or pion-pion [4] systems in Eq. (2) are described in terms of the corresponding form factors. The D -wave meson-meson form factors are represented by relativistic Breit-Wigner formulae. In the transition form factors for $R_{S,P,D}^{M_2M_3}$ we choose $R_S^{K^0\pi} \equiv K_0^*(1430)$, $R_P^{K^0\pi} \equiv K^*(892)$, $R_S^{\pi\pi} \equiv f_0(980)$ and $R_P^{\pi\pi} \equiv \rho(770)^0$.

A selected amplitude From Eqs. (1) and (3) one obtains for the $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ S -wave annihilation amplitude with $[\pi^+\pi^-]_S$ subsystem in the final state,

$$A_{n \ 2S} = -\frac{G_F}{2} A_1 \chi_2 a_2 f_{D^0}(m_K^2 - s_0) F_0^{K^0 f_0}(m_{D^0}^2) F_0^{\pi^+\pi^-}(s_0), \quad (4)$$

where $s_\pm = (p_{\pi^\pm} + p_{K^0})^2$, $s_0 = (p_{\pi^+} + p_{\pi^-})^2$. Other amplitudes can be derived similarly, their expression will be given in a forthcoming paper, however some explicit formulae can be found in Ref. [6].

3 Experimental $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ data

Isobar model and unitarity The experimental Dalitz plot analyses [1, 2] are performed within the isobar model to describe the final state meson-meson interactions and many free parameters are used (of the order of 2 per amplitude): the BABAR Collaboration




Fig. 1 Result of the present fit compared to the Dalitz-plot projection of the Belle data [2]

model relies on 43 parameters and that of Belle on 40. Amplitudes are not unitary neither in the 3-body channels nor in the 2-body sub-channels. Two-body unitarity is incorporated in the present model: unitary form factors are used in the $K\pi$ S - and P -wave [7] and in the $\pi\pi$ S -wave [4] amplitudes. The branching fraction of the sum of these amplitudes is larger than 80% of the total $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ branching fraction.

4 Preliminary results and concluding remarks

The present model has 28 free parameters, mostly unknown transition form factors and a minimization procedure is used to reproduce the $K_S^0 \pi^-$, $K_S^0 \pi^+$ and $\pi^+ \pi^-$ squared effective mass $m_{\pm}^2 = s_{\pm}$ and $m_0^2 = s_0$ projections of the experimental Dalitz plot analyses [1,2]. Preliminary results are shown in Fig. 1 for the Belle Collaboration analysis [2]. Similar results are obtained with the BABAR analysis [1].

Conclusion This theoretically constrained analysis might be useful to improve the determination of the D^0 - $\bar{D}^0$ mixing parameters and of the CKM angle γ .

References

1. Del Amo Sanchez, P., *et al.* (BABAR Collaboration): Measurement of $D^0 - \bar{D}^0$ Mixing Parameters Using $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $D^0 \rightarrow K_S^0 K^+ K^-$ Decays. Phys. Rev. Lett. 105, 081803 (2010)
2. Zhang, L.M., *et al.* (Belle Collaboration): Measurement of $D^0 - \bar{D}^0$ Mixing Parameters in $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ Decays. Phys. Rev. Lett. 99, 131803 (2007)
3. Beneke, M., Neubert, M.: QCD factorization for $B \rightarrow PP$ and $B \rightarrow PV$ decays. Nucl. Phys. B 675, 333 (2003)
4. Dedonder, J.-P., Furman, A., Kamiński, R., Leśniak, L., Loiseau, B.: Final state interactions and CP violation in $B^+ \rightarrow \pi^+ \pi^- \pi^\pm$ decays. Acta Phys. Pol. B 42, 2013 (2011)
5. Barton, G.: Dispersion Techniques in Field Theory, W. A. Benjamin, INC., New York (1965)
6. Leśniak, L., *et al.*: Dalitz plot studies of $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. To appear in Meson2012 proceedings, E. P. J. Web of Conferences, <http://www.epj.org/>, arXiv:1209.4805v1 [hep-ph]
7. El-Bennich, B., *et al.*: CP violation and kaon-pion interactions in $B \rightarrow K \pi^+ \pi^-$ decays. Phys. Rev. D 79, 094005 (2009)